

**This Page Is Inserted by IFW Operations  
and is not a part of the Official Record**

## **BEST AVAILABLE IMAGES**

**Defective images within this document are accurate representations of the original documents submitted by the applicant.**

**Defects in the images may include (but are not limited to):**

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORLED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C07D 239/30</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/29166</b> <b>(43) International Publication Date:</b> 2 November 1995 (02.11.95)
<b>(21) International Application Number:</b> PCT/GB95/00676 <b>(22) International Filing Date:</b> 27 March 1995 (27.03.95) <b>(30) Priority Data:</b> 9408270.8 26 April 1994 (26.04.94) GB <b>(71) Applicant (for all designated States except US):</b> ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BOWDEN, Martin, Charles [GB/GB]; Carr Green Close, Rastrick, Brighouse, West Yorkshire HD6 3LX (GB). BROWN, Stephen, Martin [GB/GB]; 4 Dearnfield, Upper Cumberworth, Huddersfield HD8 8NX (GB). JONES, John, David [GB/GB]; 57 Keats Road, Greenmount, Burry, Lancashire BL8 4ED (GB). <b>(74) Agents:</b> TIERNEY, Francis, John et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealott's Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG12 6YA (GB).		<b>(81) Designated States:</b> BR, CA, CN, CZ, HU, JP, KP, PL, RO, RU, SK, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PROCESS FOR THE PREPARATION OF 4,6-DICHLOROPYRIMIDINE  <b>(57) Abstract</b>  A process for preparing 4,6-dichloropyrimidine comprising treating 4,6-dihydroxypyrimidine with phosgene in the presence of a suitable base and optionally in the presence of a solvent.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## PROCESS FOR THE PREPARATION OF 4,6-DICHLOROPYRIMIDINE

The present invention relates to a process for converting 4,6-dihydroxypyrimidine (1) into 4,6-dichloropyrimidine (2) using phosgene and a suitable base. 4,6-Dichloropyrimidine is useful as a chemical intermediate in the agrochemical industry. It is especially useful in the preparation of ICIA5504.

It is known, for example, that phosphoryl chloride in the presence of dimethylaniline will convert 4,6-dihydroxypyrimidine to 4,6-dichloropyrimidine (Journal Chemical Society (1943) 574-5, and *ibid* (1951) 2214). The problem with these methods is that, when used on a large scale, a considerable amount of phosphoric acid by-product is produced and this must be disposed of in some way.

The present invention provides a process for preparing 4,6-dichloropyrimidine comprising treating 4,6-dihydroxypyrimidine with phosgene in the presence of a suitable base.

4,6-Dihydroxypyrimidine (1) can also exist in the tautomeric forms (A) and (B) and references to 4,6-dihydroxypyrimidine include all its tautomeric forms.

Suitable bases include tertiary amines of formula  $R^1R^2R^3N$  (wherein  $R^1$ ,  $R^2$  and  $R^3$  are, independently,  $C_{1-10}$  alkyl, aryl, heteroaryl or aryl( $C_{1-4}$ )alkyl) and heterocyclic amines optionally substituted by  $C_{1-10}$  alkyl. Examples of tertiary amines are triethylamine, 4-(*N,N*-dimethylamino)-pyridine, *N,N*-diisopropylethylamine and especially dimethylaniline. Examples of heterocyclic amines are pyridine, 2-methylpyridine, 4-methylpyridine, imidazole and *N*-alkyl pyrrolidines (such as *N*-methylpyrrolidine).

It is preferred that the base:phosgene molar ratio is in the range 1:10 to 10:1, especially in the range 1:1 to 1:4 (such as 2:3 and 2:4.5).

Alkyl groups are straight or branched chain and, unless stated otherwise, preferably contain from 1 to 6, especially from 1 to 4, carbon atoms. Examples are methyl, ethyl, *iso*-propyl, *n*-propyl, *n*-butyl and *tert*-butyl.

Aryl is preferably phenyl.

Heterocyclic amines are preferably saturated or unsaturated 3-7 membered carbon nitrogen rings. They are, for example, pyridine,

- 2 -

imidazole, pyrrolidine or piperidine.

Heteroaryl is a 3-7 membered carbon nitrogen ring. It is, for example, pyridine, imidazole, pyrazole or pyrrolidine.

It is preferred that the process is carried out in a solvent or mixture of solvents. Chlorinated solvents (such as dichloromethane, 1,1,2,2-tetrachloroethane or chlorobenzene), ethers (such as tetrahydrofuran, glyme, diglyme or triglyme), polar aprotic solvents [such as esters (for example a C<sub>1-4</sub> alkyl ester, such as methyl formate or iso-propyl formate) or nitriles (for example propionitrile, butyronitrile, benzonitrile or acetonitrile)] are preferred. Mixtures of solvents include, for example, a mixture of acetonitrile and dichloromethane.

The process is preferably carried out in the temperature range -10°C to 130°C, especially 0°C to 120°C, particularly 10°C to 90°C.

In one aspect the present invention provides a process for preparing 4,6-dichloropyrimidine comprising adding phosgene to a mixture of 4,6-dihydroxypyrimidine and a suitable base.

In a further aspect the present invention provides a process for preparing 4,6-dichloropyrimidine comprising adding phosgene to a mixture of 4,6-dihydroxypyrimidine and a suitable base, wherein all the phosgene to be used in the process is added at the beginning of the process.

In another aspect the present invention provides a process for preparing 4,6-dichloropyrimidine comprising adding phosgene to a mixture of 4,6-dihydroxypyrimidine and a suitable base (such as dimethylaniline or diisopropylethylamine) in a chlorinated solvent, wherein the molar ratio of 4,6-dihydroxypyrimidine:suitable base:phosgene is in the range 1:(0.8 to 2.5):(2.5 to 3.6) especially in the range 1:(1.5 to 2.2):(2.9 to 3.3).

In yet another aspect the present invention provides a process for preparing 4,6-dichloropyrimidine comprising adding phosgene to a mixture of 4,6-dihydroxypyrimidine and a suitable base (such as dimethylaniline or diisopropylethylamine) in a nitrile solvent, wherein the molar ratio of 4,6-dihydroxypyrimidine:suitable base:phosgene is in the range 1:(0.1 to 2.4):(4 to 9) especially in the range 1:(0.1 to 2.1):(4.4 to 6.5).

In a further aspect the present invention provides a process for preparing 4,6-dichloropyrimidine the process comprising adding phosgene to a mixture of 4,6-dihydroxypyrimidine and a suitable base (such as dimethylaniline) in a suitable solvent (such as dichloromethane,

- 3 -

acetonitrile or tetrahydrofuran), all the phosgene to be used in the process being added at the beginning of the process, and heating the reaction mixture (preferably for 1 to 30, especially 1 to 6 or 15 to 24 hours). The suitable base can be recovered (in salt form) during the product isolation and can be recycled.

The following Examples illustrate the invention. The apparatus used in the following Examples was dried before use, and reactions were conducted under nitrogen using anhydrous conditions.

#### EXAMPLE 1

4,6-Dihydroxypyrimidine (0.94g) was suspended in dichloromethane, dimethylaniline (1.12g) added and phosgene (5g) was then condensed into the mixture. The resulting mixture was heated at reflux for 24 hours, then cooled and poured into water. High pressure liquid chromatographic (hplc) analysis of the resulting organic layer showed a 4,6-dihydroxypyrimidine: 4,6-dichloropyrimidine ratio of 39:58.

#### EXAMPLE 2

4,6-Dihydroxypyrimidine (20.5g) was dispersed with agitation in dichloromethane (400ml). Dimethylaniline (40.4g) was added to the agitated mixture and the system was sealed (except for a vent line to a scrubber). Phosgene gas (56g) was introduced from a cylinder and condensed onto a cold finger and collected in a pressure equalised dropping funnel. Once collected, the phosgene liquid was added to the reaction mixture over 15 minutes. The mixture was heated and agitated at reflux (29°C approximately) for 17 hours after which time the mixture was cooled to room temperature and the excess phosgene removed by sparging with nitrogen.

Water (400ml) was added slowly to the agitated reaction mass with cooling to maintain the temperature at ambient. The organic layer was separated, and the aqueous was then extracted with dichloromethane (2 x 100ml). The combined extracts were dried over anhydrous sodium sulphate and concentrated by rotary evaporation to give 4,6-dichloropyrimidine as an orange crystalline solid (27g), equivalent to a yield of 80% (hplc analysis).

#### EXAMPLE 3

4,6-Dihydroxypyrimidine (2.0g) was dispersed with agitation in acetonitrile (40ml), dimethylaniline (2.1g) was added and the mixture was heated to 50°C. Phosgene gas (14.6g) was added to the mixture (by bubbling

- 4 -

it through the mixture) over 1 hour. The mixture was kept at 50°C for 4.5 hours, cooled to room temperature, and the excess phosgene removed by sparging with nitrogen. Analysis (hplc) of the resulting reaction mass showed it to comprise 4,6-dichloropyrimidine (in 81% yield).

#### EXAMPLE 4

To a mixture of 4,6-dihydroxypyrimidine (5.14g, 1 equivalent) and imidazole (6.19g, 2 equivalents) in acetonitrile (100ml) was added phosgene (28g, 6.2 equivalents). The resulting mixture was stirred for 2½ hours at room temperature and for 1 hour at 50°C. The reaction mixture was purged with nitrogen overnight and then partitioned between water and dichloromethane. The organic layer was separated and the aqueous extracted twice more with dichloromethane. The organic extracts were combined, washed with water (twice), dried over magnesium sulphate and evaporated to dryness to leave 4,6-dichloropyrimidine as a pale yellow solid.

#### EXAMPLE 5

To a stirred mixture of 4,6-dihydroxypyrimidine (5.18g, 1 equivalent) and 4-(N,N-dimethylamino)pyridine (0.55g, 0.1 equivalent) in acetonitrile (100ml) was added phosgene (28g, 19.7ml, 6.2 equivalents) in two aliquots. The resulting mixture was stirred for 10 minutes at room temperature and was then stirred for 4 hours at 55°C. The reaction mixture was purged with air after which water (200ml) was added. The resulting mixture was extracted with dichloromethane (3x100ml). The organic extracts were combined, washed with water (100ml), dried over magnesium sulphate and evaporated to dryness to leave 4,6-dichloropyrimidine (4.63g).


#### EXAMPLE 6

To a stirred mixture of 4,6-dihydroxypyrimidine (5.18g, 1 equivalent) and N,N-diisopropylethylamine (11.75g, 2 equivalents) in acetonitrile (100ml) was added phosgene (28g, 19.7ml, 6.2 equivalents) in two aliquots. The resulting mixture was stirred for 10 minutes at room temperature and was then stirred for 4 hours at 55°C. The reaction mixture was sparged with air overnight after which water (100ml) was added. The resulting mixture was extracted with dichloromethane (3x100ml). The organic extracts were combined, washed with water (100ml), dried over magnesium sulphate and evaporated to dryness to leave 4,6-dichloropyrimidine (6.35g).


- 5 -

## CHEMICAL FORMULAE



(In Description)



4,6-dichloropyrimidine



4,6-dihydroxypyrimidine

 $\text{COCl}_2$  phosgene $\text{POCl}_3$  phosphoryl chloride



- 6 -

CLAIMS

1. A process for preparing 4,6-dichloropyrimidine comprising treating 4,6-dihydroxypyrimidine with phosgene in the presence of a suitable base.
2. A process as claimed in claim 1 wherein the base:phosgene molar ratio is in the range 1:10 to 10:1.
3. A process as claimed in claim 1 or 2 wherein the process is carried out in a solvent or mixture of solvents.
4. A process as claimed in claim 1, 2 or 3 wherein the base is a tertiary amine of formula  $R^1R^2R^3N$  (wherein  $R^1$ ,  $R^2$  and  $R^3$  are, independently  $C_{1-10}$  alkyl, aryl, heteroaryl or aryl( $C_{1-4}$ )alkyl) or a heterocyclic amine optionally substituted by  $C_{1-10}$  alkyl.
5. A process as claimed in claim 1, 2, 3 or 4 wherein phosgene is added to a mixture of 4,6-dihydroxypyrimidine and a suitable base.
6. A process as claimed in claim 1, 2, 3, 4 or 5 wherein the process comprises adding phosgene to a mixture of 4,6-dihydroxypyrimidine and a suitable base in a chlorinated solvent, wherein the molar ratio of 4,6-dihydroxypyrimidine:suitable base:phosgene is in the range 1:(0.8 to 2.5):(2.5 to 3.6).
7. A process as claimed in claim 1, 2, 3, 4 or 5 wherein the process comprises adding phosgene to a mixture of 4,6-dihydroxypyrimidine and a suitable base in a nitrile solvent, wherein the molar ratio of 4,6-dihydroxypyrimidine:suitable base:phosgene is in the range 1:(0.1 to 2.4):(4 to 9).
8. A process as claimed in any one of the preceding claims wherein the base is dimethylaniline or diisopropylethylamine.

## INTERNATIONAL SEARCH REPORT

 Interna I Application No  
 PCT/GB 95/00676

 A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C07D239/30

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)  
 IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	FR,A,1 310 810 (BASF) 22 October 1962 see page 8; claims; example 29	1-7
Y	EP,A,0 095 637 (BAYER) 7 December 1983 see page 1 - page 8	1-7
Y	EP,A,0 173 191 (BAYER) 5 March 1986 see page 1 - page 8; claims	1-7



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 May 1995

Date of mailing of the international search report

23. 05. 95

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+ 31-70) 340-3016

Authorized officer

Francois, J

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat'l Application No

PCT/GB 95/00676

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR-A-1310810	13-03-63	DE-B- 1164418 US-A- 3365452	23-01-68
EP-A-0095637	07-12-83	DE-A- 3220105	01-12-83
EP-A-0173191	05-03-86	DE-A- 3431698 JP-A- 61060672 US-A- 4668788	13-03-86 28-03-86 26-05-87